

Lecture 5 Completion of a discrete valuation ring

Let $(R, \mathfrak{m})$ be a discrete valuation ring and let $t \in \mathfrak{m}$ be a local parameter.

We have a group homomorphism

$$\begin{aligned} F^\times &\longrightarrow \mathbb{Z} \\ x &\longmapsto v(x) \end{aligned}$$

and extend it to a map

$$F \longrightarrow \mathbb{Z} \cup \{\infty\}$$

by letting $v(0) = \infty$. This map is known as the valuation map and it is such that for all $x, y \in F$:

$$v(x+y) \leq \min \{v(x), v(y)\}.$$

So if we fix any $c \in \mathbb{R}_{>1}$, then we get a map

$$|\cdot|_v: F \longrightarrow \mathbb{R}_{\geq 0}$$

by letting

$$x \mapsto |x|_v := \begin{cases} c^{-v(x)} & , \text{ if } x \neq 0 \\ 0 & , \text{ if } x = 0 \end{cases}$$

This map is an absolute value map as it is s.t.

$$(A1) \quad \forall x \in F: \quad |x|_v = 0 \iff x = 0$$

$$(A2) \quad \forall x, y \in F: \quad |xy|_v = |x|_v |y|_v$$

$$(A3) \quad \forall x, y \in F: \quad |x + y|_v \leq |x|_v + |y|_v$$

In fact it satisfies a property stronger than (A3), namely

$$(A3') \forall x, y \in F: |x + y|_v \leq \max\{|x|_v, |y|_v\}.$$

So we may say that it is a *nonarchimedean* absolute value. We may recover the local ring structure $(R, \mathfrak{m})$ from the absolute value. Indeed, we have

$$R = \{x \in F \mid |x|_v \leq 1\},$$

$$\mathfrak{m} = \{x \in F \mid |x|_v < 1\}.$$

All DVR's may be obtained from nonarchimedean absolute values.

As $|\cdot|_v$ is an absolute value, the map $d_v(x, y) := |x - y|_v$ turns F into a metric space. Indeed, we have

$$(M1) \quad \forall x, y \in F : d_v(x, y) = 0 \iff x = y,$$

$$(M2) \quad \forall x, y \in F : d_v(x, y) = d_v(y, x),$$

$$(M3) \quad \forall x, y, z \in F : d_v(x, z) \leq d_v(x, y) + d_v(y, z).$$

Recall that a metric space (M, d) is complete if every Cauchy sequence $\{x_i\}_{i=1}^{\infty}$ there is $x \in M$ s.t.

$$\lim_{i \rightarrow \infty} x_i = x$$

The completion $\overline{M}$ of M is an isometry $\gamma: M \rightarrow \overline{M}$ that satisfies the following universal property. Given any isometry $f: M \rightarrow C$, where C is complete, there is a unique isometry $\hat{f}: \overline{M} \rightarrow C$ that makes the diagram

$$\begin{array}{ccc} & \hat{f} & \\ & \nearrow & \\ \overline{M} & \xrightarrow{\quad} & C \\ \uparrow & \nearrow f & \\ K & & \\ M & & \end{array}$$

commute. If $M = F$ and $d(\cdot, \cdot)$ comes from an absolute value, then

$$\overline{F} = \mathcal{C}_F / \mathcal{N}_F,$$

where $\mathcal{C}_F$ is the ring of Cauchy sequences $\{x_i\}_{i=1}^{\infty}$ with ring structure given component-wise, and $\mathfrak{N}_F \subseteq \mathcal{C}_F$ is the maximal ideal that consists of the sequences $\{x_i\}_{i=1}^{\infty}$ s.t.

$$\lim_{i \rightarrow \infty} x_i = 0.$$